DIAGNOSTIC MARKER

DISEASES

ANA, ENA, C1q, Centromere B, dsDNA, Histone, Jo-1, Nucleosome, Rib-P, RNP/ Sm, RNP-70, PM-Scl-100, Scl-70, Sm, SS-A, SS-A-52, SS-A-60, SS-B, ssDNA, DNAse activity, Rheumatoid Factor, ACPA, MCV, CCP, alpha-Fodrin, Annexin V, beta-2-Glycoprotein I, Cardiolipin, Phosphatidic Acid, Phosphatidyl Inositol, Phosphatidyl Serine, Prothrombin, ANCA, BPI, Cathepsin G, Elastase, GBM, Lactoferrin, Lysozym, MPO, PR3, AMA-M2, ASCA, ASMA, SLA, LC1, LKM-1, gp210, Sp100, tTG, Gliadin, DGP, Intrinsic Factor, Parietal Cell, Calprotectin (stool), Lactoferrin (stool), TG, TPO

- Rheumatoid arthritis (RA)
- Felty's syndrome
- Juvenile chronic arthritis
- Sjoegren's syndrome
- Systemic lupus erythematosus (SLE)
- Lupus nephritis
- Neonatal lupus eryrthematosus
- Congenital heart block (CHB)
- Subacute cutaneous lupus erythemtosus (SCLE)
- Drug-induced lupus
- Systemic scleroderma
- Raynaud's syndrome
- CREST syndrome
- Polymyositis / dermatomyositis
- Idiopathic myositis
- Mixed connctive tissue disease (MCTD)
- Undifferentiated connective tissue disaese (UCTD)
- Anti-synthetase syndrome
- Cryoglobulinemia
- Hypocomplementemic urticarial vasculitis syndrome (HUVS)
- Antiphospholipid syndrome (APS)
- Systemic vasculitis
- Granulomatosis with polyangiitis (GPA)
- Microscopic polyangiitis (MPA)
- Goodpasture syndrome
- Celiac disease
- Pernicious anaemia
- Inflammatory bowel diseases
- Autoimmune hepatitis type I (AIH I)
- Autoimmune hepatitis type II (AIH II)
- Primary biliary cirrhosis (PBC)
- Primary sclerosing cholangitis (PSC)
- Hashimoto thyroiditis

relevant marker for the disease

component of diagnostic criteria